سیستون®

Himalaya HERBAL HEALTHCARE

ترکیبات:

هر قرص روکشدار سیستون حاوی مواد موثره:

نام علمی گیاه	نام انگلیسی	مقدار
عصاره‌ها		
Didymocarpus pedicellata	-	130mg
Saxifraga ligulata	-	98mg
Rubia cordifolia	Indian madder	32mg
Cyperus scariosus	Umbrella's edge	32mg
Achyranthes aspera	Prickly chaff flower	32mg
Onosma bracteatum	Sedge	32mg
Vernonia cinerea	Ash-Colored fleabane	32mg
پودرها		
-	Shilajeet (purified)	26mg
-	Lime silicate calx	32mg

مواد جانبی:
میکروکریستالین سلولز، سدیم کربوکسی متیل سلولز، منیزیم استئارات، کراس پویدون، سیلیکون دی اکسید کلوییدال، اپادرای ۲۰۰

شکل دارویی: قرص روکشدار

گروه درمانی: حل کننده سنگ (لیتوتریپتیک: تجزیه کننده سنگ کلیه و مثانه)، ضد عفونی کننده مجاری ادراری

مکانیسم عمل:
سیستون از اشباع شدن عوامل سنگ ساز جلوگیری می کند و جذب اگزامید از روده را کنترل میکند. (اگزامید ماده‌ای است که در فرمولاسیون سنگ‌های رسوبی یافت می‌شود.) و بین کریستالوئید و کلوئید تعادل ایجاد می کند. سیستون تشکیل سنگ را از طریق کاهش عوامل سنگ ساز مانند اگزالیک اسید و کلسیم هیدروکسی پرولین مهار می کند و منجر به دفع آنها از طریق پودر کردن می‌گردد. سیستون باعث متلاشی شدن سنگها و کریستال‌ها از طریق اثر بر روی موسین که ذرات را به یکدیگر پیوند می‌دهد، می‌شود. فعالیت ضد میکروبی آن درمهار عفونت‌های مجاری ادراری هنگام دفع ادراری سنگ‌ها و کریستالها بسیار مفید می‌باشد. فعالیت ضد اسپاسمی و ضد التهابی آن التهاب حالب و علائم درد و سوزش را در هنگام ادرار کردن کاهش می‌دهد.

موارد مصرف:
۱ـ سنگ‌های ادراری:
- سنگ‌های اگزالات
- سنگ‌های فسفات
- سنگ‌های اورات و اوریک اسید
- سنگ‌های کریستالی
- جلوگیری از بازگشت مجدد سنگ‌های ادراری

۲ـ یک عامل کمکی در:
- عفونت‌های مجاری ادراری
- عفونت‌های مجاری ادراری در دوران بارداری
- التهاب نامشخص پیشابراه
- التهاب حاد مثانه، التهاب لگنچه کلیه
- سوزش ادراری
- التهاب مفاصل، نقرس (از طریق ترشح اسید اوریک در ادرار)

۳ـ کاهش خروجی ادرار (به عنوان مدر عمل می کند)
۴ـ در وضعیت‌های اورژانس بی‌اختیاری ادراری در بانوان
۵ـ نقص عملکرد کلیه به دنبال التهاب کلیه
۶ـ سنگ‌های بزاقی

منع مصرف: موردی گزارش نشده است.

مقدار مصرف:

در سنگ‌ها و کریستال‌های ادراری:
۲عدد قرص روزی ۲بار به مدت ۴تا ۶ ماه و یا تا زمانیکه سنگ دفع شود.

در عفونت‌های مجاری ادراری برگشت‌پذیر:
۱عدد قرص روزی ۲بار به مدت ۶ تا ۱۲ هفته

درد حاد کلیوی:
۲عدد قرص روزی ۲بار تا زمانیکه علائم فروکش کنند.

در سوزش ادراری:
۲عدد قرص روزی ۲بار برای ۴روز تا ۲هفته یا تا زمانیکه علائم فروکش کنند.

جلوگیری از بازگشت مجدد سنگ‌های ادراری:
۱عدد قرص ۲بار در روز به مدت ۴تا ۵ ماه بعد از دفع شدن سنگ‌ها

راه مصرف: خوراکی

احتیاط و هشدارها:
سیستون در جایی که سنگ به اندازه‌ای بزرگ است که باعث انسداد می‌شود و یا در حالات اورژانسی، قابل توصیه نمی‌باشد. در کودکان با دیورز زیاد باید مانیتورینگ انجام شود.

عوارض جانبی: موردی گزارش نشده است.

مصرف در بارداری و شیردهی:
اگر چه زنان باردار و شیرده از سیستون برای مدت طولانی استفاده نموده‌اند بدون آنکه گزارشی در مورد عوارض جانبی وجود داشته باشد اما توصیه می‌شود سیستون در دوران بارداری و شیردهی فقط با مشورت پزشک مصرف گردد.

مصرف در کودکان و افراد سالمند:
کودکان و افراد سالمند می‌توانند بدون مشکلی مصرف نمایند. در کودکان با دیورز زیاد باید مانیتورینگ انجام شود.

تداخل با سایر داروها: موردی گزارش نشده است.

شرایط نگهداری:
- در دمای کمتر از ۳۰ درجه سانتی‌گراد و دور از نور مستقیم نگهداری نمایید.
- از قرار دادن فرآورده در یخچال خودداری نمایید.
- دور از دسترس کودکان نگهداری نمایید.

بسته بندی: قوطی حاوی ۶۰ قرص روکشدار

این فرآورده یک دارو است.
- این فرآورده فقط با ارائه نسخه پزشک به فروش می‌رسد.
- دارو، فراورده‌ای است که بر روی سلامتی شما تاثیر می گذارد و استفاده از آن برخلاف دستور مصرف می‌تواند برای شما خطرناک باشد.
- نحوه تجویز پزشک و روش استفاده و مصرف را از داروساز درخواست نمایید.
- پزشک و داروساز در زمینه قطع دارو و منفعت و مضرات آنها تجربه لازم را دارند.
- خودسرانه مصرف دارو را قطع ننمایید و دوره درمان را کامل کنید.
- بدون مشورت با پزشک خود نسخه را تجدید ننمایید.

تاریخ بروزرسانی آبان ۱۴۰۵

دارنده پروانه ساخت : شرکت اهران تجارت ـ ایران
تحت لیسانس شرکت هیمالایا ـ هند

Cystone®

Himalaya HERBAL HEALTHCARE

Composition of each Cystone F.C Tablet:

Extracts:		
Didymocarpus pedicellata	-	130mg
Saxifraga ligulata	-	98mg
Rubia cordifolia	Indian madder	32mg
Cyperus scariosus	Umbrella's edge	32mg
Achyranthes aspera	Prickly chaff flower	32mg
Onosma bracteatum	Sedge	32mg
Vernonia cinerea	Ash-colored fleabane	32mg
Powders:		
-	Shilajeet (purified)	26mg
-	Lime silicate calx	32mg

Inactive ingredients:
Microcrystalline cellulose, Sodium carboxymethyl cellulose, Magnesium stearate, Crospovidone, Colloidal silicon dioxide, Opadry 200.

Description:
Round standard convex transparent film coated tablets.

Pharmaceutical form: F.C Tablet

Pharmacotherapeutic group:
Lithotryptic (disintegrates renal and urinary calculi), Urinary antiseptic.

Mode of action:
Cystone prevents supersaturation of lithogenic substances, controls oxamide absorption from the intestine (a substance that precipitates stone formation) and corrects crystalloid-colloid imbalance. It inhibits calculogenesis by reducing the stone forming substances like oxalic acid, calcium hydroxyproline etc., and causes their expulsion by micropulverization. Cystone causes disintegration of the calculi and the crystals by acting on the mucin, which binds the particles together. Its antimicrobial activity is beneficial in the prevention of urinary tract infections associated with urinary stones and crystalluria. Its antispasmodic and anti-inflammatory activities relieve ureteric colic and alleviate symptoms of painful and burning micturition.

Indications:
1- Urolithiasis:
- Oxalate stones
- Phosphate stones
- Uric acid and urate calculi
- Crystalluria
- Prevents post-operative recurrence of calculi

2- As an adjuvant in:
- Urinary tract infections
- Urinary tract infections during pregnancy
- Non-specific urethritis
- Cystitis, Pyelitis
- Burning micturition
- Arthritic conditions, gout (due to its uricosuric action)

3- Decreased urine output - as a diuretic
4- Urgency, incontinence of urine in women
5- Kidney dysfunction secondary to nephritis
6- Sialolithiasis

Contraindications: None

Dosage:

In urolithiasis and crystalluria:
2 tablets twice daily for 4 to 6 months or till the stone passes out.

In recurrent urinary tract infections:
1 tablet twice daily for 6 to 12 weeks.

In acute renal colic:
2 tablets twice daily till symptoms subside.

In burning micturition:
2 tablets twice daily for 4 days to 2 weeks or till symptoms subside.

To prevent recurrence of renal calculi:
1 tablet twice daily for 4 to 5 months after the calculus is flushed out.

Route of administration: Oral

Precautions: None

Adverse effects: None

Usage during pregnancy and lactation:
Although pregnant and lactating women have used Cystone for a long time without any reported adverse effects, it is recommended that Cystone be taken under constant medical supervision during pregnancy and lactation.

Usage by children and elderly:
Can be used safely. Monitored use is recommended in children with excessive diuresis.

Interactions with other drugs: None

Special warnings:
Cystone is not advisable where the calculus size is big enough to cause obstruction, or in an emergency. Monitored use is recommended in children with excessive diuresis.

Undesirable effects: None

Storage conditions: Store below 30°C, away from direct sunlight. Keep out of the reach of children. Do not refrigerate.

Presentation: Sealed packs of 60 F.C Tablets.

Conditions for sale: The product can be sold only against a medical prescription.

THIS IS A MEDICAMENT
- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not interrupt by yourself, the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Under license of: Himalaya Wellness Company
Makali, Bengaluru 562 162 (india)
www.himalayawellness.com